

# Factor Maps

Kai Prince

2025-08-14

## Table of contents

Backlinks	1
Outlinks	1

## Backlinks

- [Group](#)
- [Measure Theory and Ergodic Theory](#)
- [Følner sequence](#)
- [Amenable](#)
- [Actions](#)

**Definition 0.1** (Kra et al. [\(2022\)](#), Definition 2.1). For a system  $(X, \mu, T)$  we say that the system  $(Y, \nu, S)$  is a *measurable factor* of  $(X, \mu, T)$  if there is a measurable map  $\pi : X \rightarrow Y$ , the *measurable factor map*, such that  $\pi(\mu) = \nu$  and  $(S \circ \pi)(x) = (\pi \circ T)(x)$  for  $\mu$ -almost every  $x \in X$ .

**Proposition 0.1** (cf. Host [\(2019\)](#), Proposition 5). Let  $(X, \Gamma)$  be a topological dynamical system where  $\Gamma$  is an amenable group,  $x_0 \in X$ , and  $\mu$  be an ergodic invariant probability measure supported on the closed orbit of  $x_0$  under the action of  $\Gamma$ .

Let  $(Z, m_Z, H)$  be the Kronecker factor of  $(X, \mu, \Gamma)$ , with factor map  $\pi : X \rightarrow Z$ .

Let  $X \times Z$  be endowed with the group action of  $\Gamma \times H$ . Let  $\tilde{\mu}$  be the measure on  $X \times Z$  and image of  $\mu$  under the map  $X \rightarrow X \times Z$  where  $x \mapsto (x, \pi(x))$ .

Then there exists a Følner sequence  $\tilde{\Phi}$  and a point  $z_0 \in Z$  such that  $(x_0, z_0)$  is generic for  $\tilde{\mu}$  along  $\tilde{\Phi}$ .

## Outlinks

- [A Short Proof of a Generalised Conjecture of Erdős for Amenable Groups](#)

- Kra, B., et al. (2022). '*Infinite sumsets in sets with positive density*', Available at: <https://arxiv.org/abs/2206.01786>.
- Host, B. (2019). '*A short proof of a conjecture of erdős proved by moreira, richter and robertson*', Available at: <https://arxiv.org/abs/1904.09952>.